



COPY OF PAPERS
ORIGINALLY FILED

Goteb.6482

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

APPLICANT: Srinivas Uppugunduri GROUP: 1646
SERIAL NO: 09/986,522 EXAMINER: B.Y. Kwon
FILED: November 2, 2001
FOR: NOVEL, SPECIFIC INHIBITORS OF ACUTE AND CHRONIC
INFLAMMATION

Assistant Commissioner of Patents
Washington, D.C. 20231

Sir:

TRANSMITTAL OF CERTIFIED COPY

Attached please find the certified copy of the foreign application from which priority is claimed for this case:

Country: Sweden
Appln No.: 9901615-6
Filing Date: 05/05/1999

Respectfully submitted,

Arlene J. Powers
Registration No. 33,298
Samuels, Gauthier & Stevens
225 Franklin Street
Boston, Massachusetts 02110
Telephone: (617) 426-9180
Extension: 110

I hereby certify that this paper (along with any paper referred to as being attached or enclosed) is being deposited with the United States Postal Service on the date shown below with sufficient postage as first class mail in an envelope addressed to the: Commissioner of Patents and Trademarks, Washington, D.C. 20231.

Emily C. Porell

Date

02/07/2003

PRV

PATENT- OCH REGISTRERINGSVERKET
Patentavdelningen

Intyg Certificate



Härmed intygas att bifogade kopior överensstämmer med de handlingar som ursprungligen ingivits till Patent- och registreringsverket i nedannämnda ansökan.

This is to certify that the annexed is a true copy of the documents as originally filed with the Patent- and Registration Office in connection with the following patent application.

(71) Sökande Srinivas Uppugunduri, Linköping SE
Applicant (s)

(21) Patentansökningsnummer 9901615-6
Patent application number

(86) Ingivningsdatum 1999-05-05
Date of filing

Stockholm, 2002-10-31

För Patent- och registreringsverket
For the Patent- and Registration Office

Hjördis Segerlund
Hjördis Segerlund

Avgift
Fee 170:-

**PATENT- OCH
REGISTRERINGSVERKET
SWEDEN**

Postadress/Adress
Box 5055
S-102 42 STOCKHOLM

Telefon/Phone
+46 8 782 25 00
Vx 08-782 25 00

Telex
17978
PATOREG S

Telefax
+46 8 666 02 86
08-666 02 86

TITLE

NOVEL, SPECIFIC INHIBITORS OF ACUTE AND CHRONIC INFLAMMATION

DESCRIPTION

5 Technical field

The present invention relates to novel, specific inhibitors of acute and chronic inflammation, method for treatment of acute and chronic inflammation and/or problems in hemostasis related to platelet function..

- 10 The object of the present invention is to obtain novel, specific inhibitors of acute and chronic inflammations in order to be able to treat such inflammations.

Background of the invention

- Three families of cell-adhesion molecules (CAMs) have been implicated in mediating
15 interactions of platelets, endothelial cells, and leukocytes: the selectins, the integrins and the immunoglobulin superfamily. The selectin family of molecules comprising E-, L-, and P-selectin act in concert with other cell adhesion molecules to effect adhesive interactions of platelets, endothelial cells, and leukocytes. Extensive literature is available implicating cell adhesion molecules in diverse disease processes including reperfusion injury, cancer,
20 coronary heart disease, atherosclerosis, restenosis after coronary angioplasty, and chronic inflammatory diseases like asthma and IBD.

- The emigration of white blood cells to inflammatory sites requires at least four steps: leukocyte rolling along activated endothelium, leukocyte activation, firm adhesion and
25 transendothelial migration. Interaction of the selectins with their carbohydrate ligands seems to be important for the initial binding of the leukocytes to the endothelium under conditions of fluid shear stress. Subsequent firm adhesion and extravasation seems to be mediated by another family of molecules, the β 2 (CD 18)-integrins. A number of soluble mediators like IL-1 β , TNF, endotoxin, thrombin and histamine can up regulate one or more endothelial
30 adhesion molecules. The major adhesion receptors and ligands regulating leukocyte-endothelium interactions include ICAM-1, ICAM-2/LFA-1; VCAM/VLA-4; L-selectin/GlyCAM-1; CD 34, MAdCAM-1; E-selectin/ESL-1, PSGL-1; P-selectin/PSGL1.

Genetic experiments involving construction and testing of mice deficient in each of these

selectins suggest that they provide overlapping, but crucial contributions to leukocyte recruitment in inflammation. Mice lacking each of these selectins exhibit defects in neutrophil rolling and extravasation. Further, mice deficient in both E and P selectin present extreme leukocytosis, elevated cytokine levels and alterations in haematopoiesis.

5

Platelet P-selectin may also play a very important role in both hemostasis and the ensuing inflammatory reaction. Platelets are rapidly recruited to the site of the vascular injury to prevent excessive bleeding. The interaction of platelets with the vessel wall is a crucial event leading to the formation of a hemostatic plug. The role of platelet P-selectin in hemostasis
10 was confirmed in a recent study which showed that platelets roll on stimulated endothelium expressing P-selectin in vivo. Furthermore, bleeding time was prolonged by 40% in P-selectin deficient mice.

Today the treatment of inflammatory conditions include treatment using steroids which ne-
15 gatively affect the immuno defence system and leads to a total inhibition of the inflammation, where it is of interest and importance to control the inflammatory regulating system.

Rational behind random in vitro screening

20 Since initial binding of leukocytes to endothelium initiates the inflammatory process, it has been utilised in an vitro system to study the adhesion of various cells to human umbilical cord endothelial cells. The method is well established and has been successfully utilised by a number of research groups. This in vitro static adhesion assay has been used herein for random screening of substances that can block leukocyte adhesion to endothelial cells. A
25 variety of substances were tested as literature has shown that a number of, both, carbohydrate and non-carbohydrate structures can block interaction of selectins with cognate ligands. Substances that can block interaction of endothelial adhesion molecules with cognate ligands on leukocytes have potential as novel, selective inhibitors of acute and chronic inflammatory reactions and ischemia reperfusion states, without risk of general
30 immunosuppression.

Description of the present invention

Random screening of various chemical entities resulted in identification of three lead compounds that could block adhesion of neutrophils. The compounds obtained in this

screening were : isomaltitol, uridine and 4-thiouridine.

The invention will be described more in detail in the following with reference to screening tests performed. The test results are also illustrated in the accompanying graphs.

5

The results of the screening of substances that can block adhesion of neutrophils to L-cells transfected with E-selectin

L-cells permanently transfected with full-length E-selectin by electroporation. Human neutrophils were isolated from EDTA whole blood by density centrifugation. Purity and
10 total number were determined by standard haematology instruments. L-cells were grown in 96-well microtiter plates.

L-cells and neutrophils were pre-incubated with respective substances for 45 min at 37°C . Neutrophils were then allowed to adhere to L-cells in the presence of different substances for
15 20 min at 37°C. Non-adherent cells were removed by repeated washings and the number of adherent cells determined by measuring myeloperoxidase content. Since binding of neutrophils to selectin is calcium dependent, EGTA (which chelates calcium) was used as a negative control.

20 All three substances inhibit adhesion of neutrophils, but to different extents (Fig 1-3). EGTA consistently blocks adhesion of neutrophils. There is no clear dose-response for these substances which could partly depend on the fact that this is a static adhesion assay and selectin mediated binding seems to function best under flow conditions. Maximal inhibition seemed to occur already at micromolar levels.

25

Random screening of substances that can block adhesion of colon cancer cells, Colo 201, to L-cells transfected with E-selectin

Although protein ligands have been characterised for the selectins, they recognise a number of carbohydrate and non-carbohydrate ligands. The hypothesis is that these compounds
30 should block E-selectin-SLe^x interaction. To confirm this it was also tested if these substances can block adhesion of Colo 201 cells to L-cells. This interaction has been shown to be mediated predominantly by E-selectin - SLe^a group of molecules.

None of these substances could inhibit binding of Colo 201 cells to HUVEC suggesting that

this is a specific inhibition of neutrophils (Fig 4-6). Once again EGTA blocked adhesion of Colo 201 cells. In contrast to previous experiments with neutrophils, a monoclonal antibody against E-selectin was also quite effective in blocking adhesion of Colo 201 cells to TNF stimulated HUVEC.

5

Inhibition of neutrophil adhesion to TNF stimulated HUVEC

To further extend the finding from L-cells the binding of neutrophils to TNF activated endothelium (HUVEC) was studied.

- 10 Endothelial cells were grown in 96-well microtiter plates to confluence. They were stimulated with TNF for 4 h at 37°C. Isolated neutrophils were incubated for 30 min at 37°C with respective substances. The endothelial cells were washed after 4 h and 2×10^5 neutrophils were added to each well and the cells were allowed to adhere in the presence of the respective substances.

15

Isomaltitol inhibited the adhesion of neutrophils to TNF stimulated HUVEC by roughly 30%. Once again a clear dose-response was not observed. In fact, the opposite seems to be true with increasing inhibition with decreasing dose (Fig 7).

- 20 4-Thiouridine was more potent in inhibiting adhesion of neutrophils to TNF stimulated HUVEC. 4-Thiouridine reduced the adhesion of neutrophils by 60 %. Maximal adhesion seemed to occur already in the micromolar levels. Fig 8 shows a representative experiment and Fig 9 summarises data from two separate experiments performed in quadruplicate. Pre-incubation of neutrophils with EGTA or monoclonal antibodies against E-selectin blocked
25 adhesion of neutrophils.

Effect of 4-Thiouridine on up-regulation of E-selectin

- The substances mentioned above should block the interaction of E-selectin with it's ligand and should not have any effect on cytokine induced up-regulation of adhesion molecules. A
30 cell-ELISA was performed to assess the surface expression of adhesion molecules.

Endothelial cells were grown in microtiter plates, stimulated with TNF or IL-1 with or without respective substances for 4 h at 37°C. After stimulation the cells were washed and fixed with formaldehyde. A cell-ELISA was performed in analogy to regular ELISA using

primary and secondary antibodies.

Simultaneous incubation of endothelium with different concentration of 4-thiouridine did not affect TNF induced up-regulation of E-selectin as judged by Cell-ELISA (Fig 10). These results suggest that 4-thiouridine inhibits neutrophil adhesion by interfering with interaction of E-selectin with cognate ligands and does not seem to affect up-regulation of E-selectin.

Viability

E-selectin transfected L-cells were incubated with 1 mM, 0.1 mM, 0.001 mM, 0.1 μ M uridine, thiouridine, or isomaltitol for 2 hours. Viability after incubation was > 90 %. Similarly neutrophils were incubated with 1 mM, 0.1 mM, 0.001 mM, 0.1 μ M uridine, thiouridine, or isomaltitol for 2 hours. Viability after incubation was 80-97%.

Potency of lead compounds

Based on data from the above experiments, the following order can be proposed 4-thiouridine > isomaltitol > uridine.

The compounds of the present invention can be administered in the form of oral, rectal, injection, or inhalatory preparations. Oral compositions normally exist as tablets, granules, capsules (soft or hard), or powders, either coated or uncoated products. As coated products they may be merely enteric coated to provide for a more readily administered preparation, or as a sustained release coated composition, where the release of active compound will take place due to the dissolution of the coating, which dissolution is dependent on where in the gastro-intestinal tract one will have a release. Thus the release can be controlled as to place and time. It may also be advantageous to coat the active compound if this is subject to degradation, such as by gastric acid, in order then to have the compound to pass the stomach.

Tablets and capsules normally contain one dose of the active compound, i.e., the dose determined to fulfill the requirements of obtaining a therapeutically active level in serum or otherwise, either this is required once, twice or more times a day (24 hrs).

Rectal compositions are normally prepared as suppositories, where the active compound is dissolved or dispersed in a waxy compound or fat having a melting temperature in the range of the body temperature, as to release the active compound when administered rectally.

Preparations for injection are commonly made for subcutaneous, intramuscular, intravenous, or intra peritoneal administration. Injection solutions are normally provided with an adjuvant to facilitate absorption of the active compound.

- 5 Preparations for inhalation are commonly present as powders which are administered either in pressurized containers with a dosing nozzle, or in an inhaler system where the powder is dosed in the system and then the patient is inhaling air through the apparatus to such degree that the powder becomes airborne and enters the respiratory tract, including the lungs. Inhalation preparation are normally used for inflammatory conditions in the respiratory tract
- 10 including the lungs.

- The compositions contain 0.5 to 99 % by weight of active compound, and the remainder is different inert, non-therapeutically active compounds which facilitate administration, preparation such as granulation, tableting, or storage. Such inert materials may, however,
- 15 have a administratively positive effect.

- The active compounds of the invention are administered in an amount of 1 to 100 mg per kilogram body weight depending on the condition of the patient, route of administration, age and body weight of the patient, and other considerations made by the physician. The most
- 20 important aspect hereby is the serum concentration which may be 0.1 to 100 mM of active compounds, in accordance with the present findings.

FIGURE LEGENDS

- 5
10
15
- FIG. 1 to 3** Adhesion of granulocytes to E-selectin transfected L-cells incubated with different concentrations of uridine, isomaltitol, and 4-thiouridine, respectively.
- FIG. 4 to 6** Adhesion of Colo 201 to E-selectin transfected L-cells incubated with different concentrations of uridine, isomaltitol, and 4-thiouridine, respectively.
- FIG. 7 to 9** Adhesion of granulocytes to TNF stimulated HUVEC co-incubated with different concentrations of isomaltitol, 4-thiouridine, and uridine, respectively
- FIG. 10** Expression of E-selectin on stimulated HUVEC co-incubated with different concentrations of 4-thiouridine.

CLAIMS

1. The use of one or more of the compounds of the group consisting of 4-thiouridine, isomaltitol, and uridine in the preparation of therapeutically effective compositions against acute or chronic inflammations, and/or problems in hemostasis related to platelet function.
5
2. The use according to claim 1, wherein the therapeutically effective compound is 4-thiouridine.
3. The use according to claim 1, wherein the therapeutically effective compound is
10 isomaltitol.
4. The use according to claim 1, wherein the therapeutically effective compound is uridine.
5. The use according one or more of claims 1-4, wherein the therapeutically effective dose is
15 1 to 100 mg per kg bodyweight.
6. Method for treatment of acute or chronic inflammations, and/or problems in hemostasis related to platelet function,
characterized in that a therapeutically effective amount of one or more the of compounds of
20 the group consisting of 4-thiouridine, isomaltitol, and uridine in the preparation of therapeutically effective compositions against acute or chronic inflammations.
7. Method according to claim 6, wherein the active compound is administered in such an amount that the serum concentration thereof is 0.1 to 100 mM.

ABSTRACT

The present invention relates to a the use of one or more of the compounds of the group consisting of 4-thiouridine, isomaltitol, and uridine in the preparation of therapeutically effective compositions against acute or chronic inflammations, and/or problems in

- 5 hemostasis related to platelet function, as well as a method for treatment of acute or chronic inflammations, and/or problems in hemostasis related to platelet function.

PRV 99-05-08 M

Adhesion of granulocytes to E-selectin transfected L-cells incubated with different concentrations of URIDINE




FIG. 1

PRV 99-05-05 M

Adhesion of granulocytes to E-selectin transfected L-cells incubated with different concentrations of ISOMALTITOL




FIG. 2

Adhesion of granulocytes to E-selectin transfected L-cells incubated with different concentrations of 4-THIOURIDINE




FIG. 3

0000000000

PRV 99-05-03 M

Adhesion of Colo 201 to E-selectin transfected L-cells incubated with different concentrations of ISOMALTITOL




FIG. 5

Adhesion of Colo 201 to E-selectin transfected L-cells incubated with different concentrations of 4-THIOURIDINE




FIG. 6

PMN 88 1014




FIG. 7

PMN 88 1014

Adhesion of granulocytes to TNF stimulated HUVEC co-incubated with 4-Thiouridine




FIG. 8

Adhesion of granulocytes to TNF stimulated HUVEC co-incubated with 4-Thiouridine




FIG. 9

Expression of E-selectin on stimulated HUVEC co-incubated with different concentration of 4-Thiouridine




FIG. 10